

# Increasing antibiotic resistance amongst Uropathogenic *E. coli* isolated from inpatients with symptomatic UTI - a three year study

Verma Y.<sup>1</sup>, Shah K.<sup>2</sup>, Hemachander S.S.<sup>3</sup>

<sup>1</sup>Dr. Yogita Verma, Assistant Professor, <sup>2</sup>Dr. Krunal Shah, Associate Professor, <sup>3</sup>Dr. S. S. Hemachander, Microbiology, Professor & Head, All the authors are attached with Department of Microbiology, SBKS Medical Institute and Research Centre, Vadodara, Gujrat, India.

**Address for Correspondence:** Dr. Krunal Shah, A-2, Staff Quarters, Sumandeep Vidyapeeth University, Pipariya, Vadodara, E-mail: krunal2012@gmail.com

## Abstract

**Introduction:** *Escherichia coli* (E.coli) is one of the most common uropathogen worldwide. In the past few years, uropathogenic E.coli has become increasingly resistant to many drugs. **Materials and Methods:** This prospective observational study was conducted from January 2014 to December 2016. Semiquantitative urine culture was done on blood agar and MacConkey agar using standard technique. Cultures showing a significant growth of 10<sup>5</sup> CFU/ml or more were further processed. Identification and analysis of antibiotic susceptibility patterns was done by VITEK<sup>®</sup>2 (bioMerieux, USA). **Results:** A total of 674 *E. coli* samples were included in the final analysis. The isolates showed highest overall sensitivity to imipenem (75.82%) and nitrofurantoin (72.25%) whereas highest resistance was observed in case of amoxicillin-clavulanic acid (84.12%) and cephalosporins (71.96-83.23%), followed by ciprofloxacin (72.70%), amikacin (64.39%) and TMP-SMX (64.24%). Over three years duration, a trend of increasing resistance was observed in the case of cephalosporins, ciprofloxacin and trimethoprim-sulfamethoxazole (TMP-SMX). Percentage of MDR *E.coli* also increased over three years. **Conclusion:** There is an increase in antibiotic resistance amongst uropathogenic *Escherichia coli* over the past three years. Routine surveillance of resistance patterns should continue in the hospitals and antibiotic policy should be constantly updated. Judicious use of antibiotics is the need of the hour.

**Keywords:** Uropathogenic *E. coli*, Increasing resistance, MDR, urinary tract infections

## Introduction

Uropathogenic *Escherichia coli* (UPEC) is the most common etiological agent responsible for uncomplicated urinary tract infections (UTI) [1]. It causes urinary tract infections in community as well as hospital settings. It is estimated that nearly 40% of women and 12% of men will experience a symptomatic UTI in their lifetime and about 25% of these women will experience a recurrence within 6 to 12 months [2]. UTI in the hospital setting is a very common phenomenon owing to long hospital stays, catheterisation, procedures performed and drug pressure. A lot of monetary and personnel resource is spent each year in treating the infection. In the past few years, uropathogenic *E.coli* have become increasingly

resistant to many drugs [3,4]. These include oral drugs like TMP-SMX, nitrofurantoin, ciprofloxacin, oral cephalosporins as well as injectables like cephalosporins, aminoglycosides and carbapenems. Hospital acquired strains are known to be more resistant than community acquired ones [5,6].

It is important to understand the changing trends of antibiotic susceptibility of isolates, in order to better modify the empirical therapy used.

The main objective of this study was to determine antibiotic susceptibility profile of *E.coli* causing UTI in the hospital setting along with monitoring the trend of antibiotic susceptibility over three years. This study presents a three year comparison of the same from a tertiary care hospital in western India.

Manuscript received: 10<sup>th</sup> August 2017  
Reviewed: 20<sup>th</sup> August 2017  
Author Corrected: 28<sup>th</sup> August 2017  
Accepted for Publication: 4<sup>th</sup> September 2017

## Material and Methods

**Study design:** This prospective observational study was conducted at our institute from January 2014 to December 2016.

**Setting:** The institute where study was conducted is a tertiary care centre and caters to patients on both outpatient and inpatient basis. The hospital is 1200 bedded.

**Participants:** Urine samples from patients admitted to the hospital and having symptomatic urinary tract infection were collected and processed.

**Inclusion criteria:** Inpatients of both sexes and more than 18 years of age having symptomatic urinary tract infection were included in the study.

**Exclusion criteria:** Patients less than 18 years of age, outpatients, patients on immunosuppressive drugs, HIV positive and patients unwilling to participate in the study were excluded from the study

**Data source:** Prospective data collection was done from the culture and antibiotic susceptibility results obtained in the microbiology laboratory at our centre.

**Study size:** A total of 2376 Urine samples were received in microbiology laboratory for urine culture in the three year study duration.

## Results

**Baseline characteristics:** A total of 2376 samples sent for urine culture were processed in the study period. 1287 (54.16%) showed significant growth ( $10^5$  CFU/ml) and 57 samples showed mixed growth.

*Escherichia coli* was isolated from 674/1287 (52.36%) samples and was the most common uropathogen isolated.

These isolates were included in the final analysis.

The number of *E. coli* isolates obtained were 280, 237 and 157 in the year 2014, 2015 and 2016 respectively.

**Antibiotic susceptibility patterns:** The antibiotic susceptibility of *E. coli* to the following drugs was included in the final analysis: amoxicillin- clavulanic acid, ciprofloxacin, cefepime, ceftriaxone, cefuroxime, nitrofurantoin, trimethoprim- sulfamethoxazole (TMP-SMX), amikacin, gentamicin and imipenem.

The overall antibiotic susceptibility patterns of the Uropathogenic *E. coli* isolated in three years are presented in Table 1.

The isolates showed highest overall sensitivity to imipenem (75.82%) and nitrofurantoin (72.25%) whereas highest resistance was observed in case of amoxicillin- clavulanic acid (84.12%) and cephalosporins (71.96-83.23%), followed by ciprofloxacin (72.70%), amikacin (64.39%) and TMP-SMX (64.24%) in the three year duration.

**Culture:** Semi quantitative urine culture was done on Blood agar and MacConkey agar using standard technique. Cultures showing a significant growth of  $10^5$  CFU/ml or more were further processed. Cultures with mixed growth on the culture media were excluded.

**Identification and Antibiotic Susceptibility Testing:** Identification and analysis of Antibiotic susceptibility patterns was done by VITEK<sup>®</sup>2 (bioMerieux, USA), as per the manufacturer's instructions.

Culture results with isolation of *E. coli* were included in the final analysis. Repeated isolates of *E. coli* from the same patient were excluded.

**Multi Drug Resistant *Escherichia coli*:** Isolates resistant to three or more classes of antibiotics were considered as multi drug resistant (MDR).

**Statistical methods:** Statistical analysis of the test results was done using SPSS (version 20.0.0).

Categorical variables were compared using chi square test. A p value of <0.05 was considered to be statistically significant.

**Ethical considerations:** The study was approved by the institutional ethics committee.

**Table-1: Overall antibiotic susceptibility of Uropathogenic *Escherichia coli* isolated from 2014 - 2016 (n=674).**

S.no	Antibiotic tested	Sensitive (Percentage)	Resistant (Percentage)	Total
1	Amoxycillin- clavulanic acid	107 (15.88)	567 (84.12)	674
2	Ciprofloxacin	184 (27.30)	490 (72.70)	674
3	Cefepime	189 (28.04)	485 (71.96)	674
4	Ceftriaxone	185 (27.45)	489 (72.55)	674
5	Cefuroxime	113 (16.77)	561 (83.23)	674
6	Nitrofurantoin	487 (72.25)	187 (27.75)	674
7	TMP-SMX	241 (35.73)	433 (64.24)	674
8	Amikacin	434 (64.39)	240 (35.61)	674
9	Gentamicin	384 (56.97)	290 (43.03)	674
10	Imipenem	511 (75.82)	163 (24.18)	674

**Comparative analysis:** It was observed in the comparative analysis of the three year data that there was no significant difference in susceptibility of *E. coli* to amoxycillin- clavulanic acid, nitrofurantoin, and imipenem. However there are interesting trends to be observed in case of ciprofloxacin, cephalosporins, TMP-SMX and aminoglycosides [Table 2].

The Susceptibility to ciprofloxacin decreased over three years with 42.5%, 21.52% and 8.91% isolates being reported as sensitive in 2014, 2015 and 2016 respectively. The decrease in antibiotic susceptibility was statistically significant in 2014 vs 2015 (p value<0.001) and 2015 vs 2016 (p value<0.001).

The susceptibility to cephalosporins tested also decreased over three years. The decrease of susceptibility to cefepime was significant in 2014-2015 [45% vs 23.20%(p value<0.001)] and 2015-2016 [23.20% vs 5.10%(p value<0.001)]. The susceptibility to ceftriaxone also significantly decreased in 2014-2015[43.93% vs 21.94% (p value <0.001)] and 2015-2016[21.94% vs 6.37%(p value <0.001)]. Similarly the decline in sensitivity to cefuroxime was found to be significant in 2014-2015 [28.93% vs 9.70% (p value<0.001)] and 2015-2016 [9.70% vs 5.73%(p value<0.001)].

Sensitivity to TMP-SMX was lesser in 2015 and 2016 as compared to 2014. The decrease was significant in 2014 vs 2015[42.85% vs 28.27% (p value<0.001)] but insignificant in 2015 vs 2016. However if data from 2014 was compared to 2015 and 2016 combined, the decrease in susceptibility was statistically significant[42.85% vs 30.71%(p value<0.001)].

The difference in antibiotic susceptibility to amikacin between 2014 and 2015 as well as 2015 and 2016 was statistically insignificant. Susceptibility to gentamicin was significantly lesser in 2015 as compared to 2014(p value<0.001) where as the difference between 2015 and 2016 was insignificant. However the difference was significant when the results from 2014 were compared to 2015 and 2016 combined[64.64% vs 51.5%(p value<0.001)].

**MDR E.coli:** Isolates resistant to three or more classes of antibiotics were considered as multi drug resistant (MDR). The percentage of MDR E. coli isolated from urine samples in 2014, 2015 and 2016 were 53.92%, 64.13% and 79.61% respectively. The most common phenotype of MDR E. coli isolated was resistant to either of the cephalosporins, ciprofloxacin and TMP-SMX. The rates of resistance to two of the most commonly used drugs for UTI - ciprofloxacin and TMP-SMX, were highest in 2016 at 91.09% and 65.60% respectively and 98/157(62.42%) isolates were concomitantly resistant to both. Concurrent resistance to ciprofloxacin, TMP-SMX and nitrofurantoin was observed in 28/157(17.83%) isolates.

**Table 2: A three year comparison of the antibiotic susceptibility patterns of Uropathogenic *Escherichia coli*.**

S.No	Antibiotic tested	2014 (n=280)		2015 (n=237)		2016 (n=157)	
		S (%)	R (%)	S (%)	R (%)	S (%)	R (%)
1	Amoxicillin-clavulanic acid	41 (14.64)	239 (85.36)	29 (12.23)	208 (87.77)	37 (23.36)	120 (76.44)
2	Ciprofloxacin	119 (42.50)	161 (57.5)	51 (21.52)	186 (78.48)	14 (8.91)	143 (91.09)
3	Cefepime	126 (45%)	154 (55%)	55 (23.20)	182 (76.80)	8 (5.10)	149 (94.90)
4	Ceftriaxone	123 (43.93)	157 (56.07)	52 (21.94)	185 (78.06)	10 (6.37)	147 (93.63)
5	Cefuroxime	81 (28.93)	199 (71.07)	23 (9.70)	214 (90.30)	9 (5.73)	148 (94.27)
6	Nitrofurantoin	202 (72.14)	78 (27.86)	167 (70.46)	70 (29.54)	118 (75.15)	39 (24.85)
7	TMP-SMX	120 (42.85)	160 (57.15)	67 (28.27)	170 (71.73)	54 (34.40)	103 (65.60)
8	Amikacin	183 (65.36)	97 (34.64)	147 (62.02)	90 (37.98)	104 (66.24)	53 (33.75)
9	Gentamicin	181 (64.64)	99 (35.36)	121 (51.05)	116 (48.95)	82 (52.23)	75 (47.77)
10	Imipenem	209 (74.64)	71 (25.36)	185 (78.06)	52 (21.94)	117 (74.5)	40 (25.5)

## Discussion

Antibiotic resistance amongst uropathogenic *E. coli* is increasing every year as found in this study. The highest overall resistance was seen in case of cephalosporins, ciprofloxacin and amoxicillin-clavulanic acid.

Resistance to fluoroquinolones in uropathogenic *E. coli* is a global phenomenon now and varied resistance rates to ciprofloxacin have been reported from different geographical regions. In a study conducted on outpatients with community acquired UTI in Spain, the authors reported 14% resistance to ciprofloxacin [7]

whereas Timothy et al.[8] from Nigeria reported 31% resistance to ciprofloxacin amongst uropathogenic *E.coli* isolated from inpatients. Karaca Y et al. in 2005 [9], reported that ciprofloxacin resistance rates in *E.coli* varied from 5.2% to 27.6% in the ten year study period. Our study found the overall resistance to ciprofloxacin to be 72.70% in the three year study period. This is similar to the resistance rates reported by Indian studies in recent times [6,10]. Our study showed that the resistance to ciprofloxacin in *E.coli* increased significantly each year from 2014 to 2016. The high

resistance rates and significant increase in resistance to ciprofloxacin found in this study may be attributed to two factors: high usage of ciprofloxacin for treating various clinical conditions and use of ciprofloxacin as empirical therapy for treating UTI while awaiting culture reports. In a surveillance study from The Netherlands, a similar yearly increase in resistance to fluoroquinolones in *E.coli* was reported in 2000 with the projection of an increase in resistance further as fluoroquinolones remain the most commonly prescribed antibiotics [11].

The results show a high rate of resistance to TMP-SMX (64.2%) which is higher than those reported from other geographical regions [7], similar to the rate reported by Niranjana V et al [12] in a study conducted on inpatients in India, and lesser as compared to some published studies from India [6,10]. In our study, higher resistance rates to TMP-SMX were observed in 2015 and 2016 combined as compared to a 2014. In a 10 year study from Turkey, resistance rates to TMP-SMX varied from 38.5% to 69.3% [9]. Such variations of resistance rates amongst hospitals and varying resistance rates with time may be attributable to different/changing practices involving use of TMP-SMX for therapy.

Resistance to cephalosporins ranged from 71.96% to 83.23% in the present study which is similar to recent reports from Indian Hospitals [6,12]. World wide, cephalosporin resistance in uropathogenic *E.coli* has been reported to be in the range of 15.7% to 83% depending upon the geographical region and inpatient or outpatient setting [8,13,14,15]. Resistance to cephalosporins significantly increased over three years of study period. This may be attributable to excess use of cephalosporins in the community and hospital settings. In our hospital set up cephalosporins are very commonly used as first line empirical therapy for treating many clinical conditions. Findings of the present study highlight the hazardous effects of the same.

The percentage of MDR *E. coli* found by this study is comparable to that of recently published studies [12,14]. The number of MDR *E.coli* has steadily increased over the three year duration. Indiscriminate usage of antibiotics in the outpatient and inpatient care system is possibly the reason behind this. Many published studies have clearly established that prior antibiotic usage is a risk factor for infection with MDR *E.coli* [16,17,18,19,20].

According to Infectious diseases society of America and European society of Microbiology and infectious diseases, fluoroquinolones, TMP-SMX, nitrofurantoin and amoxicillin-clavulanic acid are the recommended first line drugs for uncomplicated UTI [21].

However the recommendations for the first line treatment should be decided according to the local susceptibility patterns of a region/ country. Recently published NCDC guidelines for antibiotic use in infectious diseases recommend nitrofurantoin/ TMP-SMX/ ciprofloxacin as first line antibiotics for treating uncomplicated and Piperacillin- Tazobactam/ amikacin for complicated UTI [22]. Beta lactam antibiotics are not the choice for empirical therapy possibly owing to their very high resistance rates as reflected in our study results with only 16% strains of *E.coli* being susceptible to amoxicillin -clavulanic acid.

Nitrofurantoin susceptibility is found to be nearly 72% in the present study and it did not vary over the three years significantly. However the declining susceptibility of ciprofloxacin is a major concern as it still remains as a first line drug. Ciprofloxacin has been recently named as one of the “alert drugs” by NCDC which means that it is among the drugs most frequently prescribed irrationally and should be used only when clearcut microbiological indications exist. Switching over to susceptible/narrow spectrum antibiotics after obtaining culture results while treating UTI is also strongly recommended by NCDC [22]. These measures might be able to reduce or stop the progression of the existing fluoroquinolone resistance.

**Strengths of the study:** This study presents a comparative analysis of susceptibility of *Escherichia coli* over three years duration. Therefore not only we are able to observe the trend of increasing antibiotic resistance but also the data will be helpful in fine tuning the antibiotic policy for empirical therapy in patients with UTI. According to the findings of this study, fluoroquinolones should be used as sparingly as possible.

**Limitations:** The data on actual antibiotic usage could not be collected. A correlation of antibiotic usage and increasing resistance patterns would have provided us with a better insight into the resistance pattern.

**Future directions:** Routine surveillance of antibiotic susceptibility patterns of uropathogenic *Escherichia*

*coli*, especially to ciprofloxacin, should continue. Indiscriminate usage of antibiotics should be strictly controlled.

## Conclusion

There is an increase in antibiotic resistance amongst uropathogenic *Escherichia coli* over the past three years. Judicious use of antibiotics is the need of the hour.

**Author contributions:** Verma Yogita was involved in data collection, data analysis, manuscript preparation and manuscript revision. Shah Krunal was involved in data collection, data analysis, manuscript revision. Hemachander SS was involved in manuscript revision.

**Sources of support:** None

**Acknowledgement:** None

**Presented at meeting:** None

## References

- Gupta K, Hooton TM, Wobbe CL, Stamm WE. The prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in young women. *Int. J. Antimicrob. Agents.* 1999 May; 11(3-4):305-308. Doi.org/10.1016/S0924-8579(99)00035-7.
- Foxman B. Recurring urinary tract infection: incidence and risk factors. *Am J Public Health.* 1990 Mar; 80 (3):331-3.
- Bours PHA, Polak R, Hoepelman AIM, Delgado E, Jarquin A, Matute J. Increasing resistance in community-acquired urinary tract infections in Latin America, five years after the implementation of national therapeutic guidelines. *Int J Infect Dis.* 2010 Sep; 14 (9): e770-4.doi: 10.1016/j.ijid.2010.02.2264.
- Gupta K, Scholes D, Stamm WE. Increasing prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in women. *JAMA* 1999;281(8):736-738. doi:10.1001/jama.281.8.736.281:736-8.
- Mandal J, Acharya NS, Buddhapriya D, Parija SC. Antibiotic resistance pattern among common bacterial uropathogens with a special reference to ciprofloxacin resistant *Escherichia coli*. *Indian J Med Res.* 2012 Nov; 136 (5): 842-849.
- Akram M, Shahid M, Khan AU. Etiology and antibiotic resistance patterns of community acquired urinary tract infections in JNMC hospital Alighrah, India. *Ann Clin Microbiol Antimicrob.* 2007 Mar 23; 6:4. DOI:10.1186/1476-0711-6-4
- Alós JI, Serrano MG, Gomez-Garces JL, Perianes J.. Antibiotic resistance of *Escherichia coli* from community-acquired urinary tract infections in relation to demographic and clinical data. *Clin Microbiol Infect.* 2005; 11(3):199-203. DOI:10.1111/j. 1469-0691. 2004. 01057.x.
- Timothy OO, Olusesan FJ, Adesola BO, Temitayo AA, David FO, Ige OO. Antibiotic resistance pattern of bacterial isolates from cases of urinary tract infections among hospitalized and out-patients at a tertiary health facility in South Western Nigeria. *Ann trop med pub health.* 2014;7(2):130-5. <http://www.atmph.org/text.asp?2014/7/2/130/146403>.
- Karaca Y, Coplu N, Gozalan A, Oncul O, Citil BE, Esen B. Co-trimoxazole and quinolone resistance in *Escherichia coli* isolated from urinary tract infections over the last 10 years. *International J Antimicrob Agents.* 2005; 26 (1):75-7. doi: 10.1016/j.ijantimicag.2005.03.012.
- Anand N, Asthana AK, Madan M, Vats AD. Antibiotic resistance pattern in *E.coli* causing community acquired Urinary tract infections. *Journal of Advance Researches in Biological Sciences.* 2014;6 (3&4): 209-12.
- Goettsch W, van Pelt W, Nagelkerke N, Hendrix MG, Buiting AG, Petit PL, Sabbe LJ, van Griethuysen AJ, de Neeling AJ. Increasing resistance to fluoroquinolones in *escherichia coli* from urinary tract infections in the netherlands. *J Antimicrob Chemother.* 2000 Aug;46(2):223-8.
- Niranjan V, Malini A. Antimicrobial resistance pattern in *Escherichia coli* causing urinary tract infection among inpatients. *Indian J Med Res.* 2014;139(6):945-8. <http://icmr.nic.in/ijmr/2014/june/0621.pdf>
- Sorlozano A, Jimenez-Pacheco A, Luna del castillo JD, Sampedro A, Martinez-Brocal A, Miranda-Casas C, Navarro-Marí JM, Gutiérrez-Fernández J. Evolution of the resistance to antibiotics of bacteria involved

in urinary tract infections: A 7-year surveillance study. *American J Inf Control*.2014;42(10):1033 - 8. doi: 10.1016/j.ajic.2014.06.013.

14. Dehbanipour R, Rastaghi S, Sedighi M, Maleki N, Faghri J. High prevalence of multidrug-resistance uropathogenic *Escherichia coli* strains, Isfahan, Iran. *J Nat Sci Biol Med*. 2016;7(1):22-6. DOI: 10.4103/0976-9668.175020

15. Ali I, Kumar N, Ahmed S, Dasti JI. Antibiotic Resistance in Uropathogenic *E. Coli* Strains Isolated from Non-Hospitalized Patients in Pakistan. *J Clin Diagn Res*. 2014;8 (9):DC01-4. doi: 10.7860/JCDR/2014/7881.4813.

16. Shariff V A AR, Shenoy M S, Yadav T, M R. The antibiotic susceptibility patterns of uropathogenic *Escherichia coli*, with special reference to the fluoroquinolones. *J Clin Diagn Res*. 2013 Jun;7(6):1027-30. doi: 10.7860/JCDR/2013/4917. 3038. Epub 2013 Jun 1.

17. van der Starre WE, van Nieuwkoop C, Paltansing S, van't Wout JW, Groeneveld GH, Becker MJ, Koster T, Wattel-Louis GH, Delfos NM, Ablij HC, Leyten EM, Blom JW, van Dissel JT. Risk factors for fluoroquinolone-resistant *Escherichia coli* in adults with community-onset febrile urinary tract infection. *J Antimicrob Chemother*. 2011 Mar; 66 (3):650-6. doi: 10.1093/jac/dkq465. Epub 2010 Dec 1.

18. Khawcharoenporn T, Vasoo S, Singh K. Urinary Tract Infections due to Multidrug-Resistant Enterobacteriaceae: Prevalence and Risk Factors in a Chicago Emergency Department. *Emerg Med Int*. 2013;2013:258517. doi: 10.1155/2013/258517. Epub 2013 Oct 31.

19. Killgore KM, March KL, Guglielmo BJ. Risk factors for community-acquired ciprofloxacin-resistant *Escherichia coli* urinary tract infection. *Annals Pharmacother*. 2004; 38:1148-52. DOI: 10.1345/aph.1D622.

20. Cohen-Nahum K, Saidel-Odes L, Riesenber K, Schlaeffer F, Borer A. Urinary tract infections caused by multi-drug resistant *Proteus mirabilis*: Risk factors and clinical outcomes. *Infection*. 2010 Feb;38(1):41-6. doi: 10.1007/s15010-009-8460-5. Epub 2009 Dec 7.

21. Gupta K, Hooton TM, Naber KG, Wullt B, Colgan R, Miller LG, Moran GJ, Nicolle LE, Raz R, Schaeffer AJ, Soper DE. International Clinical Practice Guidelines for the Treatment of Acute Uncomplicated Cystitis and Pyelonephritis in Women: A 2010 Update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clinical Infectious Diseases*. 2011;52(5):103-120. <https://doi.org/10.1093/cid/ciq257>

22. National Treatment guidelines for Antimicrobial use in Infectious Diseases. NCDC.2016.Version 1.0. [http://pbhealth.gov.in/AMR\\_guideline7001495889.pdf](http://pbhealth.gov.in/AMR_guideline7001495889.pdf)

.....  
**How to cite this article?**

Verma Y, Shah K, Hemachander S.S. Increasing antibiotic resistance amongst Uropathogenic *E. coli* isolated from inpatients with symptomatic UTI - a three year study. *Trop J Path Micro* 2017;3(3):298-304. doi:10.17511/jopm.2017.i3.12.  
.....